

Wieslab Request Form - Autoimmune Diagnostics

SEND TO: Wieslab AB P.O. Box 50117, SE-202 11 Malmö, Sweden	CONTACT diagnostic.services@svarlifescience.com	T +46 (0)40 - 53 76 60 F +46 (0)40 - 43 28 90
REQUESTING DOCTOR/CLINIC Postal address for test result report	BILL TO / INVOICE ADDRESS Only doctors, laboratories and hospital administration can be invoiced	PATIENT DATA Full name: Birth date, Identity number:
REFERENCE NUMBER / COST CENTER	GENDER Man Woman Other	SAMPLING DATE
REQUESTING DOCTOR Name: Email: Phone:	SAMPLING MATERIAL Serum EDTA-Plasma	SPECIMEN COLLECTION INFORMATION <ul style="list-style-type: none"> For autoantibody assays, blood should be collected in plain tubes (serum tubes) without additives. 3 mL serum after centrifuging 7 mL blood (1300-1800g for 10 min) is enough for approx. 15 tests. Keep samples cold until transport. Transport samples at room temperature by ordinary mail or with cold packs if long transportation (>24h). Serum samples for complement analysis should be aliquoted into two or three tubes. The serum samples and 2,5 mL EDTA-plasma (for C3d) should be frozen and shipped on dry ice. For more information about sampling please see: www.wieslab.com/diagnostic-services/sampling
Comments (Patient history etc.)	The healthcare provider submitting the sample(s) with this request form hereby confirms that the patient (or the patient's guardian or trustee, if applicable) has been informed that the samples may be retained by Wieslab AB for a period of up to 5 years for the purpose of conducting further analyses in order to make a diagnosis, and that Wieslab AB intends to retain samples for a period of up to 5 years for the purpose of the Svar Life Science AB/Wieslab AB's future development of analysis methods and its business activities. No, the patient does not give her/his consent to save the sample. The patient is currently unable to give his or her consent in relation to retention of the sample(s).	

Select test panel, on suspicion of:

Individual tests on the reverse side →

***URGENT - results within 24 hours (during laboratory hours)**

If the sample arrives no later than 10.00 on a weekday, the test is performed on the same day.

Telephone number for urgent reporting:

501	ANCA for Systemic Vasculitis/anti-GBM Goodpastures syndrome Anti-GBM (Goodpasture, GP), PR3-ANCA, Capture PR3-ANCA, MPO-ANCA, Capture MPO-ANCA, ds-DNA
040	ANCA expanded analysis ANCA-panel (BPI, Azurocidine, Cathepsin G, Elastase, Laktoferrin, Lysozyme)
100	Additional testing for ANCA - IgG-subclass determination GBM (IgG1-4) 145 GBM (IgG4)
504	ANCA follow up testing of Systemic vasculitis PR3-ANCA, Capture PR3-ANCA 505 MPO-ANCA, Capture MPO-ANCA
500	Glomerulonephritis Anti-GBM, PR3-ANCA, Capture PR3-ANCA, MPO-ANCA, Capture MPO-ANCA, anti-dsDNA, ANA screen, anti-Streptolysin O, anti-PLA2R and anti-THSD7A
584	Primary Membranous Nephropathy (PMN) anti-PLA ₂ R and anti-THSD7A (IIF screen) URGENT*
247	PMN follow up test anti-PLA ₂ R (EUSA)
510	Phospholipid autoantibodies Anti-Cardiolipin (IgG + IgM), anti-Beta-2-glycoprotein 1 (IgG + IgM)
579	Expanded analysis - Antiphospholipid syndrome (APS) anti-Cardiolipin (IgA), anti-Beta-2-glykoprotein (IgA), anti-phosphatidylserine (IgGAM)
520	Connective tissue disease – undifferentiated PR3-ANCA, capture PR3-ANCA, MPO-ANCA, capture MPO-ANCA, ANA screen, anti-Cardiolipin (IgG + IgM), anti-Beta 2 glycoprotein 1 (IgG + IgM)
536	Cystic fibrosis BPI-ANCA IgA+ IgG
531	Diabetes Islet cell antibodies, anti-GAD, anti-IA2, anti-Insulin, anti-Zinc transporter 8
530	Sudden deafness ANA screen, anti-HSP-70, PR3-ANCA, Capture PR3-ANCA
524	Congenital heart block ENA screen (incl. SSA/Ro-52)
555	Inflammatory Myopathy (Myositis) Antibodies against HMGCR, cN-1A, Mi-2α, Mi-2β, TIF1γ, MDA5, NXP2, SAE1, Ku, PM-Scl100, PM-Scl75, Jo-1, SRP, PL-7, PL-12, EJ, OJ och Ro-52. <i>Only analysed in serum.</i>
551	incl. ANA screen (panel 555 incl. ANA screen)

570	Complement activation Complement function (classical, alternative and lectin (MBL) pathways), C3, C4, C3d and C1q <i>Serum samples for complement analysis should be aliquoted into three tubes. The serum samples and EDTA-plasma (for C3d) should be frozen and shipped on dry ice.</i>
572	Complement deficiency Complement function (classical, alternative and lectin (MBL) pathways), C3, C4, C3d, C1q and properdin <i>Serum samples for complement analysis should be aliquoted into three tubes. The serum samples and EDTA-plasma (for C3d) should be frozen and shipped on dry ice.</i>
571	Hereditary angioedema (HAE) C1-inhibitor (protein and function) and C4 <i>Serum samples for complement analysis should be aliquoted into two tubes. The serum samples should be frozen and shipped on dry ice.</i>
513	Inflammatory bowel diseases (IBD) ASCA IgA+IgG, IIF ANCA If IIF-ANCA is positive, a follow-up with ANCA expanded analysis 040 is recommended. <i>(for genetic analysis, please see request form: Autoimmunity & Genetics)</i>
515	Autoimmune hepatitis ANA screen, smooth muscle antibodies (SMA), liver/kidney microsomes antibodies (LKM), mitochondrial antibodies (AMA) and liver screen (LKM-1, LC-1, AMA-M2, M2-3E (BPO), Sp100, PLM, gp210, SLA/LP, Ro-52)
575	Expanded analysis - Autoantibodies in autoimmune Pancreatitis (type 2, AIP2) Anti-Carboanhydras II, Lactoferrin <i>(for genetic analysis, please see request form: Autoimmunity & Genetics)</i>
512	Celiac disease Transglutaminase IgA, IgA quantification If IgA deficiency is determined analysis for Transglutaminase IgG, Deamidated Gliadin IgG will be performed.
578	Expanded analysis, Celiac disease (Pediatric) Endomysium IgA, Deamidated Gliadin IgA
550	Pemphigus/Pemphigoid (Bullous dermatosis) Anti-BP180, BP230, Desmoglein 1, Desmoglein 3, Envoplakin and Collagen VII <i>(not available as individual tests)</i>
522	Sjögren's syndrome/ Scleroderma/systemic sclerosis ANA screen, ENA screen
527	Expanded analysis - Scleroderma Antibodies against Fibrillarín, Th/To, RNA polymerase, PM/Scl
541	Drug induced lupus ANA screen, IIF-ANCA, anti-Histones, anti-Cardiolipin (IgG + IgM)
511	Systemic lupus erythematosus (SLE) ANA screen, anti-dsDNA, ENA screen, anti-Cardiolipin (IgG + IgM)

Mark for test

ANCA

008 IIF-ANCA (C- and P-ANCA)

COMPLEMENT ANALYSIS (SERUM + EDTA-PLASMA)

Samples for complement analysis should be frozen and shipped on dry ice.

890 Anti-C1q
925 C1-inhibitor (function)
926 C1-inhibitor (level)
920 C1q
900/910 C3, C4
930 C3d (*Note! EDTA-Plasma, frozen*)
911 C4d (*Note! EDTA-Plasma, frozen*)
573 Complement function: Classical-, alternative- and lectin- (MBL) pathways
931 Properdin

NEPHROLOGY - ANTIBODIES

001 Goodpasture (GBM)
245 Phospholipase A2 receptor (PLA₂R) Screen (IIF)
247 Phospholipase A2 receptor (PLA₂R) follow up (ELISA)
140 Thrombospondin type 1 domain-containing protein 7A receptor (THSD7A)
097 Tubular basement membrane (TBM)

ORGAN/TISSUE SPECIFIC ANTIBODIES

086 Adrenal cortex
514 ASCA IgA + IgG
850 Carboanhydras II
072 Deamidated Gliadin IgA
073 Deamidated Gliadin IgG
071 Endomysium IgA
089 GAD (Glutamat dekarboxylas) - Serum (ELISA)
111 IA-2 (Islet cell antigen 2)
110 Insulin
770 Intrinsic factor (INF)
090 Islet cell
083 Liver/kidney microsomal (LKM)
018 Lever screen (LKM-1, LC-1, AMA-M2, M2-3E (BPO), Sp100, PML, gp210, SLA/LP, Ro-52)
084 Mitochondrial (AMA)
087 Parietal cell
082 Smooth muscle (SMA)
780 Steroid 21 hydroxylase (S21HY)
085 Striated muscle
120 Total IgA quantification
074 Transglutaminase IgA
147 Transglutaminase IgG
113 Zinc transporter 8 (ZnT8)

NUCLEAR ANTIBODIES

020 ANA screen on HEp-2 cells
023 dsDNA (Crithidia)
024 ENA screen (nRNP/Sm, Sm, SS-A/Ro-52, SS-B, Scl-70, Jo-1)
600 Fibrillarin
098 Histones
030 Jo-1
026 nRNP (snRNP, U1-RNP)
031 Ribosomal RNP (rRNP, P-protein)
029 Scl-100
025 Sm
027 SSA/Ro-60
032 SSA/Ro-52
028 SSB (La)
720 Th/To

MYOPATHIES (MYOSITIS)

141 cN-1A, Cytosolic 5'-nucleotidase 1A, (Mup44, NT5c1A)
485 HMGCR (HMG-CoA reductas)
481 KS (AsparaginyI-tRNA-Synthatase)
548 Myopathy (Myositis) screen (*only analysed in serum*) (Mi-2 α , Mi-2 β , TIF1 γ , MDA5, NXP2, SAE1, Ku, PM-Scl100, PM-Scl75, Jo-1, SRP, PL-7, PL-12, EJ, OJ, Ro-52)
715 PM/Scl (p100)
450 RNA polymeras

(For ANA and ENA screen, please see Nuclear antibodies)

PHOSPHOLIPID ANTIBODIES

517 Beta 2 glycoprotein 1 IgG + IgM
146 Beta 2 glycoprotein 1 IgA
516 Cardiolipin IgG + IgM
017 Cardiolipin IgA
404 Phosphatidylserine IgA
405 Phosphatidylserine IgG
403 Phosphatidylserine IgGAM screen
406 Phosphatidylserine IgM

OTHER ANTIBODY TESTS

095 HSP-70
859 Lactoferrin
240 Streptolysin O (ASO)

Information about tests, sampling instructions as well as Terms and Conditions are available on www.svarlifescience.com/services/wieslab-diagnostic-services

The latest version of the request forms are always available for download on our website. For tests related to Neurology, Therapeutic Drug Monitoring and Autoimmune & Genetics please use respective request form.

Contact us!

Are you **looking for a collaborating partner** for analyzes in clinical trials and/or projects?

Can't find the analysis or test you are looking for on the Request Form?

Send us an email diagnostic.services@svarlifescience.com

Beställningar

Send us request forms:

Autoimmune Diagnostics

Neurology

Monitoring of Biopharmaceuticals (TDM)

Autoimmunity & Genetics