

本製品は日本国法に基づき製造販売されたものです。  
 製品および文書は日本国外の法規に準じているわけではありません。  
 本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。  
 また、必要な時に読めるように大切に保管してください。  
 This product has been manufactured and sold based on Japanese law.  
 Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.  
 This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.  
 Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure  
 Eisai.jp

Category-3 OTC Drug

For skin roughness and acne

## チョコラBBピュア

### Chocola BB Pure

Skin roughness, acne and other skin troubles have a variety of causes, such as aging, irregular lifestyle, hormone imbalances and stress, which can disturb skin metabolism. In addition to the activated form of vitamin B<sub>2</sub>, which aids fat metabolism and normalizes the turnover of skin cells, チョコラ BB ピュア contains vitamin C, which helps with the production of collagen and 3 other vitamin B groups that aid healthy skin production to alleviate skin roughness and acne from inside the body.



Package	Recommended retail price (tax included)
40 tablet bottle	¥1,009
80 tablet bottle	¥1,821
130 tablet bottle	¥2,675
170 tablet bottle	¥3,169

Related website



[Chocola.com](http://Chocola.com) (in Japanese)

### Indications

#### Relief of the following symptoms:

Skin roughness, acne, dermatitis, rashes, sores, eczema, stomatitis, angular stomatitis, canker sores, glossitis, congestion of the eyes, itchy eyes, rosacea

Please consult with a physician or pharmacist if no improvement is seen in these symptoms after a month of use.

#### Supplementation of vitamin B<sub>2</sub> in the following cases:

When physically fatigued, during pregnancy and lactation, or when physically weakened during and after illness

### Dosage and Administration

Take the following dosage orally twice a day, after breakfast and dinner, with cold or hot water.

Age	Dosage	Number of doses
Adult (15 years or older)	1 tablet	Twice a day
Child (Under 15 years)	Do not take this medicine	

## Active Ingredients and Actions

The daily adult dose of 2 tablets contains the following active ingredients.

Ingredient	Amount	Action
Riboflavin Sodium Phosphate (Vitamin B <sub>2</sub> phosphate ester)	38 mg	<ul style="list-style-type: none"><li>• Supports healthy skin and mucosa.</li><li>• Aids fat metabolism, converting it to energy.</li></ul>
Calcium ascorbate (equivalent to 103 mg of vitamin C)	125 mg	<ul style="list-style-type: none"><li>• Aids collagen production, supporting healthy skin and mucosa.</li></ul>
Thiamine Nitrate (Vitamin B <sub>1</sub> nitrate)	20 mg	<ul style="list-style-type: none"><li>• Aids carbohydrate metabolism, converting it to energy.</li></ul>
Pyridoxine Hydrochloride (Vitamin B <sub>6</sub> )	24 mg	<ul style="list-style-type: none"><li>• Supports healthy skin and mucosa.</li><li>• Aids protein metabolism, converting it to energy.</li></ul>
Nicotinamide	40 mg	<ul style="list-style-type: none"><li>• Supports healthy skin and mucosa.</li></ul>

### Inactive Ingredients

Talc, calcium carbonate, corn starch, hydroxypropyl cellulose, partial alpha starch, gum arabic, carnauba wax, titanium oxide, red ferric oxide, stearic acid, magnesium stearate, shellac, cellulose, silicon dioxide, sucrose, hypromellose, pullulan, povidone, macrogol, dibasic calcium phosphate

### Precautions for Ingredients

1. Administration of vitamin C is known to influence the results of urine and stool tests, so when undertaking urine and stool testing, please inform your physician that you are taking this medicine.
2. Taking this medicine may cause urine to turn bright yellow, but this is due to assimilation of the vitamin B<sub>2</sub> contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.



## Precautions

### Consultation

1. If the following symptoms are observed after taking this medicine, these may be adverse reactions. Immediately discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash
Digestive organs	Stomach discomfort

2. The following symptom may be observed after taking this medicine. If this symptom persists or worsens, discontinue the use of this medicine and show this informational leaflet to with a physician, pharmacist or registered salesperson for a consultation.

Diarrhea

3. If symptoms do not improve even after taking the medicine for about 1 month, stop taking the medicine and consult a physician, pharmacist or registered salesperson, bringing this informational leaflet with you.

### Precautions for Storage and Handling

1. Store in a dry, cool place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. Also, do not transfer other medications, etc., into this container. (It may lead to misuse or alter the quality.)
4. Moisture may alter the outer appearance of the tablets, so do not handle with wet hands.
5. The packing material inside the container is provided to prevent damage to the tablets during transport. After opening the cap, discard the packing material.
6. A desiccant is included in the container. Do not ingest the desiccant.
7. Do not take tablets that have exceeded their expiration date.
8. From the standpoint of maintaining quality, as a rule of thumb, once the cap has been opened, use the tablets within 6 months, regardless of their expiration date. There is a space marked "Date Opened" on the box's inner flap. Please use that to record the date of opening.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): [https://inquiry.eisai.co.jp/webapp/form/17672\\_sdab\\_2/index.do](https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do)