

本製品は日本国法に基づき製造販売されたものです。
 製品および文書は日本国外の法規に準じているわけではありません。
 本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。
 また、必要な時に読めるように大切に保管してください。
 This product has been manufactured and sold based on Japanese law.
 Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.
 This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.
 Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure
 Eisai.jp

Category-3 OTC Drug

Stomatitis care whenever and wherever needed

チョコラBB® 口内炎リペアショット

Chocola BB Repair Shot

Do you experience painful eating or difficulty speaking due to stomatitis or throat inflammation? チョコラ BB 口内炎リペアショット is a medicated spray effective against stomatitis or pain due to throat inflammation. チョコラ BB 口内炎リペアショット acts directly on the affected area through dual actions of disinfection and mucosal repair. It is a convenient, easy-to-carry spray with a cap that can be used without getting hands dirty.



| Package | Recommended retail price (tax included) |
|---------|--|
| 30 mL | ¥1,133 |

Related website



Chocola.com (in Japanese)

Indications

Stomatitis, sore throat, swollen throat, rough throat, throat discomfort or hoarse voice due to throat inflammation

Dosage and Administration

Spray an appropriate amount to the affected area several times a day.

Precautions for Dosage and Administration

1. Follow directions for dosage and administration.
2. When allowing children to use the medicine, a guardian must be present to watch and tell them how to use it.
3. Use only as a spray onto the affected area in the mouth or throat. Do not ingest.
4. If you inhale while applying, the medication may enter the bronchial tubes or lungs. Therefore, when applying it to the throat, aim the nozzle at the affected area and spray while exhaling or vocalizing.
5. Avoid contact with eyes. In the case that the medicine comes into contact with eyes, immediately rinse eyes with cold or lukewarm water, and immediately see an ophthalmologist for medical treatment.

How to use the spray

Remove the cap, bring the bottle to your mouth, direct the nozzle to the affected area, and press down on the pump head to spray.

- * When using for the first time, press the pump head repeatedly until the medication comes out.
- * When pressing the pump head, press it down firmly and completely, not part way.
- * Use several times a day, spraying 1 to 3 times on each application.

Active Ingredients and Actions

100 mL contains the following ingredients.

| Ingredient | Amount | Action |
|--------------------------------------|--------|--|
| Azulene sulfonate sodium hydrate | 20 mg | • Repairs mucous membrane, promoting healing of the affected area. |
| Cetylpyridinium chloride monohydrate | 300 mg | • Disinfects the affected part. |

Inactive Ingredients

Benzyl benzoate, ethanol, glycerin, sodium saccharin, D-sorbitol, propylene glycol, beeswax, L-menthol, dibasic sodium phosphate, flavor, sodium dihydrogen phosphate



Precautions

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur]

Do not take this medicine for a long time.

Consultation

- The following people should contact a physician, dentist, pharmacist or registered salesperson for a consultation before use.
 - People undergoing treatment from a physician or dentist
 - People who have experienced allergic symptoms associated with drugs, etc.
 - People with the following symptoms
Erosion inside the mouth
- If the following symptoms are observed after using this drug, these may be adverse reactions. Immediately discontinue the use of this medicine and show this informational leaflet to a physician, dentist, pharmacist or registered salesperson for a consultation.

| Parts Affected | Symptom |
|------------------|----------------------------|
| Skin | Rash/redness, itching |
| Mouth | Irritation |
| Digestive organs | Stomach discomfort, nausea |

- If symptoms do not improve even after using this treatment for 5 or 6 days, stop taking this medicine and consult a physician, dentist, pharmacist or registered salesperson, bringing this informational leaflet with you.

Precautions for Storage and Handling

- Store with the cap on in a cool place away from direct sunlight.
- Strong pressure applied to the sides of the bottle may result in fluid leakage.
- Keep out of reach of children.
- Do not transfer to a different container. (It may lead to misuse or alter the quality.)
- If the medication gets onto clothing, wash immediately with water and/or detergent.
- Do not take medicine that has exceeded its expiration date.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do