

chocola bb syrop

チョコラBBこどもシロップ

Category-3 OTC Drug



< チョコラBBブランド情報

製品詳細・添付文書

+

overview

Product Description

チョコラBB[®]こどもシロップ

For decreased physical strength during or after illness, stomatitis, eczema, and tiredness in little children (3 months or over).

Vitamin B₂ has been called the “vitamin for fat metabolism” and the “vitamin for skin and mucosa.” Deficiencies of this vitamin can cause symptoms such as fatigue, eczema, dermatitis and stomatitis to occur easily.

Chocola BB Children's Syrup contains vitamin B₂ as its main ingredient, along with vitamins B₁ and B₆ and biotin, which is known as vitamin for anti-dermatitis. It is a peach-flavored syrup form for children to swallow easily.

Recommended for improving symptoms, such as eczema, acne, dermatitis and stomatitis, in little children aged 3 months or over, as well as for supplementation of vitamin B₂ when physical strength is decreased during or after illness.

Indications

▽ Dosage and Administration

▽ Ingredients

▽ Precautions

▽ Packaging and Price

Indications



Relief of the following symptoms:

Skin roughness, rashes, sores, acne, eczema, dermatitis, stomatitis, angular stomatitis, canker sores, glossitis, rosacea, congestion of the eyes, itchy eyes

Please consult with a physician or pharmacist if no improvement is seen in these symptoms after a month of use.

Supplementation of vitamin B₂ in the following cases:

When physically fatigued, or when strength is decreased during and after illness

Dosage and Administration

Take twice a day after meals, using the measuring cup provided to measure the dose

Age	Dosage	Number of doses
Between 7 and 15 years	15mL	Twice per day
Between 1 and 7 years	7.5mL	Twice per day
Between 3 months and 1 year	5mL	Twice per day
Under 3 months	Do not take this medicine	-

Precautions for Dosage and Administration

When allowing children (3 months–15 years) to take the medicine, a guardian must be present to watch and tell them how to take it.

Ingredients

Active Ingredients and Actions

One 30 mL bottle contains the following active ingredients.

Ingredients	Amount	Action
Riboflavin Sodium Phosphate (Vitamin B ₂ phosphate ester)	19mg	Supports healthy skin and mucosa. - Aids fat metabolism, converting it to energy.
Pyridoxine Hydrochloride (Vitamin B ₆)	12mg	Supports healthy skin and mucosa. Aids protein metabolism, converting it to energy.
Thiamine Nitrate (Vitamin B ₁ nitrate)	6mg	Aids carbohydrate metabolism, converting it to energy.
Nicotinamide	20mg	Supports healthy skin and mucosa.
Biotin	250µg	Supports healthy skin and mucosa and suppresses inflammation of the skin.

Inactive Ingredients

Sodium benzoate, ethanol (0.1 mL or less), citric acid, sodium citrate, D-sorbitol, paraben, flavor, hydrogenated maltose starch syrup, glycine, propyl gallate, povidone, DL-malic acid, pH adjuster

Precautions for Ingredients

Taking this medicine may cause urine to turn bright yellow, but this is due to assimilation of the vitamin B₂ contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.

Precautions

▲ Precautions for Use

Consultation

1. If the following symptoms are observed after taking this drug, these may be adverse reactions. Immediately discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash
Digestive organs	Stomach discomfort

2. The following symptom may be observed after taking this medicine. If this symptom persists or worsens, discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist or registered salesperson for a consultation.
Diarrhea
3. If symptoms do not improve even after taking the medicine for about 1 month, stop taking the medicine and consult a physician, pharmacist, or registered salesperson, bringing this informational leaflet with you.

Precautions for Storage and Handling

1. Store sealed in a cool place away from direct sunlight.
2. Keep out of reach of children.
3. Keep refrigerated after opening the bottle, and use it as soon as possible.
4. Do not transfer to a different container. Also, do not transfer other medications, etc., into this container. (It may lead to misuse or alter the quality.)
5. After administration, rinse the measuring cup, and store it securely with the syrup.
6. Do not take medicine that has exceeded its expiration date.

Packaging and Price

Packaging	Recommended retail price
30mL×2	770yen, tax included (700yen, tax excluded)
30mL×2×5	3,850yen, tax included (3,500yen, tax excluded)