

onji

遠志（オンジ）の恵み

Category-3 OTC Drug



< 遠志（オンジ）の恵み情報

製品詳細・添付文書

+

overview

Product Description

オンジ 遠志の恵み®

For improvement of forgetfulness after middle age

Recommended for people who experience symptoms of forgetfulness after middle age.

- If you are pronouncing “that” and “it” more and more frequently in the conversation.
- If you say the same things repeatedly.

Recovers your original “ability to remember” by improving the “symptoms of forgetfulness” due to aging. A natural drug product containing Polygala root extract as ingredient in small and easy to take tablets, which requires 1 tablet per time (3 times daily).

* ONJI-NO-MEGUMI does not have therapeutic or preventive effects on dementia.

* Notification of revision of Precautions

Point

- Small and easy to take tablets, requiring only 1 tablet per time.
- Recovers your original “ability to remember” by improving memory-related functions.

Indications

▼ Dosage and Administration

▼ Ingredients

▼ Precautions

▼ Packaging and Price

Indications



Improves forgetfulness after middle age

scroll

Dosage and Administration

Take the following dose with cold or hot water before or between meals.

Age	Dosage	Number of doses
Adults (15 years or over)	1 tablet	3 tablets per day
Under 15 years	Do not take this medicine	-

Ingredients

Active Ingredients and Actions

The daily dose (3 tablets) contains the following ingredients.

Ingredients	Amount
Polygala root extract	536 mg (quantity of the raw herb polygala: 3 g)

Additives

Talc, caramel, carnauba wax, titanium oxide, magnesium stearate, cellulose, hypromellose, macrogol, and silicon dioxide

Precautions relating to ingredient and amount

This medicine may interfere with some test values on diabetes.

Precautions

▲ Precautions for Use

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen, or adverse reactions are more likely to occur.]

This product should not be taken by the following persons:

1. Persons diagnosed as having the following: dementia
2. Persons are suspected to have dementia

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
 1. Patients undergoing medical treatment from a physician
 2. Pregnant women or women suspected of being pregnant
 3. Patients who have experienced allergic symptoms associated with drugs, etc.
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching
Gastrointestinal system	Nausea/vomiting, loss of appetite, feeling of discomfort in the stomach
Neuropsychiatric system	Dizziness

3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Diarrhea

4. When symptoms do not improve even after taking this medicine for about 1 month, stop taking this medicine and consult a physician, pharmacist, or registered salesperson, being sure to take this document with you.
5. If you take this medicine for a long time, consult a physician, pharmacist, or registered salesperson.

Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight with the package bag zipped tightly.
2. Keep out of reach of children.
3. Do not transfer the medicine to a different container. Moreover, do not put other drugs in this container. (This may lead to misuse or alter the quality of the drug.)
4. Do not use the product after the expiration date.
5. Since humidity may change the appearance of the tablets, do not touch them with wet hands.

Packaging and Price

Packaging	Recommended retail price
30 tablets	1,518yen, tax included (1,380yen, tax excluded)
84 tablets	4,048yen, tax included (3,680yen, tax excluded)