

saclon q tablets

サクロン錠

Category-2 OTC Drug



<

サクロンブランド情報

製品詳細・添付文書

+

overview

Product Description

サクロン錠

A green stomach medicine that refreshes the stomach. For stomach retching, heartburn, and excessive drinking.

Excessive secretion of gastric acid may cause stomach retching and heartburn. As another cause, alcohol may directly irritate the stomach mucous membrane itself due to excessive drinking.

Saclon Tablets attenuate (neutralize) the action of excess gastric acid, suppresses its secretion, and restore and protect the damaged stomach mucous membrane with a green ingredient made from chlorophyll.

For people who are suffering stomach retching, gripping pain, and nausea when brushing their teeth in the morning. Take Saclon Tablets, which will give you refreshing effects.

Indications

▼ Dosage and Administration

▼ Ingredients

▼ Precautions

▼ Packaging and Price

Indications



Heartburn, excessive drinking, stomachache, hyperacidity, stomach heaviness, feeling of discomfort in the stomach, feeling of fullness in the stomach, heaviness in the stomach, heaviness in the chest, belching, nausea (retching, stomach retching, retching due to hangovers and overdrinking, sick feeling, and feeling of sickness), vomiting

Dosage and Administration

Take the following dose with cold or hot water between meals and before going to bed on an empty stomach.

Age	Dosage	Number of doses
Adults (15 years or over)	4 tablets	3 times per day
Between 8 and 15 years	2 tablets	3 times per day
Under 8 years	Do not take this medicine	-

*Between meals means, about 2-hours after meals, and when food is digested already.

Precautions for Dosage and Administration

When allowing children (between 8 and 15 years) to take the medicine, a guardian must be present to watch and tell them how to take it.

Ingredients

Active Ingredients and Actions

The daily adult dose of 12 tablets contains the following ingredients.

Ingredients	Amount	Action
Potassium copper chlorophyllin	120mg	It repairs and protects mucous membranes of stomach.
Anhydrous dibasic calcium phosphate	1,020mg	It directly neutralizes the excess gastric acid.
Precipitated calcium carbonate	1,020mg	It directly neutralizes the excess gastric acid.
Magnesium hydroxide	960mg	It directly neutralizes the excess gastric acid.
Scopolia extract	30mg	It suppresses the secretion of gastric acid and stops the pain.

Additives

Corn starch, hydroxypropyl cellulose, l-menthol, flavor, potassium chloride, cinnamon bark, magnesium stearate, povidone

Precautions for Ingredients

1. The stool may turn green after taking this drug, but this is due to excretion of the ingredient potassium copper chlorophyllin (green). It is not a cause for concern.
2. Saclon Tablets do not use aluminum-containing ingredients.

Precautions

⚠ Precautions for Use

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen, or adverse reactions are more likely to occur.]

1. This drug should not be taken together with the following drugs:
Pain relief/antispasmodic medicine for the stomach
2. Nursing women should either not take this medicine or stop nursing while taking this medicine.
(It may cause the infant's pulse to quicken if you transfer to breast milk.)

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
 1. Patients undergoing medical treatment from a physician
 2. Pregnant women or women suspected of being pregnant
 3. The elderly
 4. Patients who have experienced allergic symptoms associated with drugs, etc.
 5. Persons with the following symptoms:
Urination difficulty
 6. Persons diagnosed as having the following
Kidney disease, heart disease, glaucoma, thyroid disease
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching

3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Dry mouth, constipation, diarrhea
4. When symptoms do not improve even after taking this medicine for about 2 weeks, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Other precautions

Some may have trouble lactating.

Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. Moreover, do not put other drugs in this container.
(This may lead to misuse or alter the quality of the drug.)
4. Note the following precautions for packets of 4 tablets each.
When taking the tablets remaining in a packet divided in multiple portions, fold back the opening of the packet before storing it, and do not take the tablets stored for more than 2 days.
5. Note the following precautions for packages of 96 tablets.
 1. The padding in the container is to prevent the tablets from breaking during transport.
Dispose of it after opening the cap of the container. Seal the cap of the container tightly for storage.

2. From the standpoint of maintaining quality, as a rule of thumb, once the cap has been opened, use the tablets within 6 months, regardless of their expiration date. There is a space marked "Date Opened" on the box's inner flap. Please use that to record the date of opening.
6. Do not use the product after the expiration date.

Packaging and Price

Packaging	Recommended retail price
40 tablets	1,078yen, tax included (980yen, tax excluded)
96 tablets	2,134yen, tax included (1,940yen, tax excluded)