

travelmin cherop lemon

トラベルミン チュロップレモン  
味 (Brand name : トラベル  
ミン チュロップレモン味)



Category-2 OTC Drug

< [トラベルミンブランド情報](#)

[製品詳細・添付文書](#) +

overview

## Product Description

**トラベルミン® チュロップレモン味**

Brand name : トラベルミン チュロップレモン味

**For prevention and relief of motion sickness**

Travelmin Churopp Lemon Flavor is a medicine that prevents and relieves symptoms associated with motion sickness, such as dizziness, nausea, and headache, to help you to comfortably enjoy your trip and going out.

It is a drop type medicine, which is easy to take for children, you can take it quickly on the spot when hurrying to go out or feeling sick.

Indications

▼ Dosage and  
Administration

▼ Ingredients

▼ Precautions

▼ Packaging and  
Price

### Indications



Prevention and relief of dizziness, nausea, and headache associated with motion sickness

## Dosage and Administration

To prevent motion sickness, take the following dose 30 minutes before boarding, either by chewing the medicine or dissolving it in your mouth.

Age	Dosage	Number of doses
15 years or over	2 tablets	Up to 2 times in intervals of at least 4 hours
11 to 14 years	2 tablets	Up to 2 times in intervals of at least 4 hours
5 to 10 years	1 tablets	Up to 2 times in intervals of at least 4 hours
Less than 5 years	Do not take this medicine	-

When taking additional doses, leave an interval of at least 4 hours between doses. Do not take this medicine more than twice per day.

### Precautions for Dosage and Administration

When allowing children (5 to 14 years) to take the medicine, a guardian must be present to watch and tell them how to take it. When giving the medicine to be taken on a school trip, for example, give instructions about the dosage, method of administration, and how to take the medicine in advance.

### Note

- This medicine is taken either by chewing it or dissolving it in your mouth. If it is difficult to chew, dissolve it before taking it, so that you do not damage your teeth. Take care that it does not accidentally become stuck in your throat.
- Take 1 tablet at a time, and do not put 2 tablets in your mouth at once.

## Ingredients

### Ingredients and their actions

The dose of 2 tablets (11 years or over) contains the following ingredients.

Ingredients	Amount	Action
d-chlorpheniramine maleate	1.33 mg	Acts on the autonomic nerves, preventing and relieving the symptoms of motion sickness.
scopolamine hydrobromide hydrate	0.166 mg	Reduces sensory confusion caused by riding in vehicles.

### Additives

Citric acid, l-menthol, flavor, acesulfame potassium, isomalt, glycerol esters of fatty acids, vitamin B<sub>2</sub>

## Precautions

### ▲ Precautions for Use

#### When not to use the product

If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.

1. This drug should not be taken together with the following drugs:

Other medicine for motion sickness, cold medicine, antipyretics/analgesics, sedatives, expectorant, indigestion and heartburn relief, internal medicine containing antihistamines, etc. (such as internal medicine for rhinitis and medicine for allergies)

2. Do not operate vehicles or machinery after taking this drug. (Symptoms such as sleepiness, blurred vision, and abnormal brightness may occur.)

#### Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

1. Patients undergoing medical treatment from a physician
2. Pregnant women or women suspected of being pregnant
3. The elderly
4. Patients who have experienced allergic symptoms associated with drugs, etc.
5. Persons with urination difficulty
6. Persons diagnosed with glaucoma or heart disease

2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this box to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching
Neuropsychiatric system	Headache
Urinary system	Urination difficulty
Other	Facial hot flashes, abnormal brightness

On rare occasions, the following serious symptoms may occur. In this case, visit a physician immediately.

Name of symptom	Symptom
Aplastic anemia	Symptoms such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.
Agranulocytosis	Symptoms such as acute high fever, chills and sore throat, etc. may occur.

3. The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this drug, and show this box to your physician, pharmacist, or

registered salesperson for a consultation.

Dry mouth, constipation, drowsiness, blurred vision

#### Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. (This may lead to misuse or alter the quality of the drug.)
4. Do not use the product after the expiration date.

#### Packaging and Price

Packaging	Recommended retail price
6 tablets	521yen, tax included (474yen, tax excluded)